

CEVA ANIMAL HEALTH, LLC

301 ROUTE 17 N., 12TH FLOOR, RUTHERFORD, NJ, 07070

Toll-Free: 1-800-999-0297

Toll-Free Fax: 1-877-777-5138

Website: www.ceva-us.com

ADAPTIL DOG APPEASING PHEROMONE DIFFUSER

Ceva Animal Health

In mammals, lactating females release substances called “appeasing pheromones”, the function of which is to reassure the offspring. Canine “appeasing pheromones” are secreted by the sebaceous glands of the intermammary sulcus in the lactating bitch.

These pheromones calm the puppy during times of stress and provide reassurance, particularly in unknown environments and when encountering novel experiences.

Research has shown that the reassuring properties of these pheromones persist even into adult age.

Adaptil uses Dog Appeasing Pheromone to reproduce the properties of the natural “appeasing pheromones” of the bitch.

Composition:

Dog appeasing pheromone (D.A.P.®)	2%
Excipients to	100 g

Uses:


Adaptil helps stop or prevent fear and stress-related signs in puppies and adult dogs, such as: Destructive behavior, Excessive vocalization, House soiling, Excessive licking

Adaptil comforts the dog in stressful environments: Visitors, Strangers, Novel and unpredictable situations leading to phobic reactions

Adaptil helps establish the puppy in a new environment: Adoption, Moving, Coping with strangers

Instructions for use:

- 1 Remove the vial cap.
- 2 Screw the diffuser onto the vial and gently tighten.
- 3 Plug the diffuser into an electric socket for 1 month.



Place the Adaptil Diffuser in the room most used by the dog.

Plug diffuser unit in an open area. Do not place behind furniture.

Coverage area: 500 - 750 sq. ft. 50 to 70 m²

One vial lasts approximately four weeks.

For best results: Plug in diffuser 1 day prior to acclimating the dog to the area.

Diffuser is intended for continual use. Although fluid may remain after 6 weeks, the vial should be replaced.

Duration of use may be adapted according to the recommendations of your veterinarian.

Presentation:

Complete kit contents:

One Diffuser + One 48 ml vial

Refill:

One 48 ml vial only

The diffuser is specially designed for this product. The properties of the product cannot be guaranteed if a different device is used.

It is recommended to renew the Adaptil diffuser after 6 months of continuous use or after the use of 6 refills (including the original one). When not in use, the Adaptil diffuser should be stored in a dust-free environment, preferably in its original packaging. Keep the pack insert for future reference.

Important safety instructions

CAUTION

When using electric products, basic precautions should always be followed, including the following:

Read all instructions before using.

To reduce the risk of injury, close supervision is necessary when a product is used near children. Adult assembly required. The surfaces of the device reach high temperatures to create evaporation of active ingredient-these surfaces should not be touched when plugged in.

Keep out of reach of children.

In case of contact with eyes, wash them immediately with water and seek the advice of a physician.

Avoid contact with skin. In case of contact, wash thoroughly with soap and water. If the product is swallowed, consult a doctor immediately and show him/her the packaging.

RISK OF ELECTRICAL SHOCK

Use only in properly functioning 120-volt electrical outlet. No adaptor or electrical cord should be used with this product. Check that the house voltage is the same as is indicated on the device. This electric device should only be used with this Adaptil Diffuser formulation. During and immediately after use, do not touch the device with your hands.

Do not immerse in water or in damp/wet areas. When plugged in, do not touch the device with metal objects or wet hands.

RISK OF FIRE

Do not cover. Do not place underneath or behind furniture. Keep away from fabrics, bedding and other material that might present a fire hazard. Use of other substance may increase the toxicity or flammability of the product.

Empty packaging and any remnants of the product should be disposed of in accordance with current regulations covering disposal of waste products.

The effects of this product were not tested by Underwriters Laboratories.

Adaptil is not a veterinary medicine. If there are signs of disease consult your veterinarian.

NAC No.: 13280000