



BAYER HEALTHCARE LLC
Animal Health Division

P.O. BOX 390, SHAWNEE MISSION, KS, 66201-0390

Customer Service Tel.: 800-633-3796

Customer Service Fax: 800-344-4219

Website: www.bayer-ah.com

LEGEND® / LEGEND® MULTI DOSE



(hyaluronate sodium) Injectable Solution

LEGEND® MULTI DOSE

(hyaluronate sodium) Injectable Solution

For Intravenous Use in Horses Only. Not for Intra-Articular Use

LEGEND®

(hyaluronate sodium) Injectable Solution

4 mL For Intravenous Use In Horses Only

2 mL For Intravenous or Intra-Articular Use In Horses Only

Bayer Animal Health

CAUTION:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION:

Legend[®] (hyaluronate sodium) Injectable Solution and Legend[®] Multi Dose (hyaluronate sodium) Injectable Solution is a clear, colorless solution of low viscosity. Legend Injectable Solution is pyrogen free, sterile and does not contain a preservative. Legend Injectable Solution 2 mL is administered by intravenous or intra-articular injection. Legend 4 mL is for intravenous use only. Legend Multi Dose Injectable Solution is pyrogen free and sterile. It is administered by intravenous injection only.

Hyaluronic acid, the conjugate acid of hyaluronate sodium, is extracted from the capsule of *Streptococcus* spp. and purified, resulting in a form which is essentially free of protein and nucleic acids.

Legend Injectable Solution is supplied in 2 mL (20 mg) and 4 mL (40 mg) vials. Each mL contains 10 mg hyaluronate sodium, 8.5 mg sodium chloride, 0.223 mg sodium phosphate dibasic, and 0.04 mg sodium phosphate monobasic. The pH is adjusted to between 6.5 and 8.0 with sodium hydroxide or hydrochloric acid.

Legend Multi Dose Injectable Solution is supplied in 20 mL vials. Each mL contains 10 mg hyaluronate sodium, 8.5 mg sodium chloride, 0.223 mg sodium phosphate dibasic, 0.04 mg sodium phosphate monobasic and 15.63 mg benzyl alcohol as a preservative. The pH is adjusted to between 6.5 and 8.0 with sodium hydroxide or hydrochloric acid.

CHEMISTRY:

Hyaluronic acid, a glycosaminoglycan, can exist in the following forms depending upon the chemical environment in which it is found: as the acid, hyaluronic acid; as the sodium salt, sodium hyaluronate (hyaluronate sodium); or as the hyaluronate anion. These terms may be used interchangeably but in all cases, reference is made to the glycosaminoglycan composed of repeating subunits of D-glucuronic acid and N-acetyl-D-glucosamine linked together by glycosidic bonds. Since this product originates from a microbial source, there is no potential for contamination with dermatan or chondroitin sulfate or any other glycosaminoglycan.

CLINICAL PHARMACOLOGY:

Hyaluronic acid is a naturally occurring substance present in connective tissue, skin, vitreous humour and the umbilical cord in all mammals. High concentrations of hyaluronic acid are also found in the synovial fluid. It also constitutes the major component of the capsule of certain microorganisms. The hyaluronic acid produced by bacteria is of the same structure and configuration as that found in mammals.

The actual mechanism of action for hyaluronate sodium in the healing of degenerative joint disease is not completely understood. One major function appears to be the regulation of normal cellular constituents. This effect decreases the impact of exudation, enzyme release and subsequent degradation of joint integrity. Additionally, hyaluronate sodium exerts an anti-inflammatory action by inhibiting the movement of granulocytes and macrophages.¹

Hyaluronate molecules are long chains which form a filter network interspersed with normal cellular fluids. It is widely accepted that injection directly into the joint pouch enhances the healing of inflamed synovium by restoring lubrication of the joint fluid. This further supplements the visco-elastic properties of normal joint fluid.

INDICATIONS:

Legend[®] (hyaluronate sodium) Injectable Solution and Legend[®] Multi Dose (hyaluronate sodium) Injectable Solution is indicated in the treatment of joint dysfunction of the carpus or fetlock in horses due to non-infectious synovitis associated with equine osteoarthritis.

DOSAGE AND ADMINISTRATION:

Legend Multi Dose (20 mL) and Legend 4 mL (40 mg) injected intravenously only. Legend 2 mL (20 mg) injected intravenously or intra-articularly in the carpus or fetlock.

Treatment may be repeated at weekly intervals for a total of three treatments.

Legend Injectable Solution 2 mL: Strict aseptic technique should be observed when administering by intraarticular injection. As with any intra-articular procedure, proper injection site disinfection and animal restraint are important. Excess joint fluid should be aseptically removed prior to injection. Care should be taken to avoid scratching the cartilage surface with the tip of the injection needle. Diffuse swelling lasting 24 to 48 hours may result from movement of the needle while in the joint space.

For intravenous administration, use aseptic technique and inject slowly into the jugular vein.

Horses should be given stall rest after treatment before gradually resuming normal activity.

Legend Injectable Solution 2 mL and 4 mL only: Discard any unused portion of the drug and the empty vial after opening.

CONTRAINDICATIONS:

There are no known contraindications for the use of Legend® (hyaluronate sodium) Solution and Legend® Multi Dose (hyaluronate sodium) Injectable Solution in horses.

RESIDUE WARNING:

Do not use in horses intended for human consumption.

HUMAN WARNINGS:

Not for use in humans. Keep this and all other drugs out of reach of children.

ANIMAL SAFETY WARNING:

For 4 mL only - Not for Intra-articular use.

Multi Dose not for Intra-articular use. The Intra-articular safety of hyaluronate sodium with benzyl alcohol has not been evaluated.

PRECAUTIONS:

Radiographic evaluation should be carried out in cases of acute lameness to ensure that the joint is free from serious fractures. As with any intra-articular (Legend Injectable Solution - 2 mL only) treatment, special precautions must be followed as to injection technique and sterility for prevention of possible swelling or infection. Intra-articular injections should not be made through skin that has been recently fired or blistered, or that has excessive scurf and counterirritant on it.

The safety of Legend Injectable Solution and Legend Multi Dose Injectable Solution has not been evaluated in breeding stallions or in breeding, pregnant or lactating mares.

ADVERSE REACTIONS:

No local or systemic side effects were observed in the Legend Injectable Solution clinical field trials with either intravenous or intra-articular injections.

Post-Approval Experience: While all adverse reactions are not reported, the following adverse reactions are based on voluntary post-approval reporting for Legend Injectable Solution:

Following **intravenous** use: Occasional depression, lethargy, and fever.

Following **intra-articular** (Legend Injectable Solution - 2 mL only) use: lameness, joint effusion, joint or injection site swelling, and joint pain.

For medical emergencies or to report adverse reactions, call 1-800-422-9874.

EFFECTIVENESS:

Effectiveness studies utilizing Legend Multi Dose Injectable Solution were not performed. Legend Multi Dose Injectable Solution was approved based on the conclusion that the effectiveness of Legend Multi Dose Injectable Solution will not differ from that demonstrated for the original formulation of Legend Injectable Solution.

Forty-six horses with lameness in either the carpal or fetlock joints were treated intravenously or intraarticularly with Legend Injectable Solution in a well-controlled clinical study conducted at four locations. One, two or three injections were given based on clinical improvement. Overall clinical improvement was judged as excellent or good in 90% of the cases treated intravenously and 96% of those treated intra-articularly with Legend Injectable Solution.

ANIMAL SAFETY:

Animal safety studies utilizing Legend Multi Dose Injectable Solution were not performed. Legend Multi Dose Injectable Solution was approved based on the conclusion that the safety of Legend Multi Dose Injectable Solution will not differ from that demonstrated for the original formulation of Legend Injectable Solution.

Legend Injectable Solution was administered to normal horses at one, three and five times the recommended intra-articular dosage of 20 mg and the intravenous dosage of 40 mg. Treatments were given once weekly for nine consecutive weeks (three times the maximum duration). No systemic clinical signs were observed nor were there any adverse effects upon hematology or clinical chemistry parameters. A transient, slight to mild post-injection swelling of the joint capsule occurred in some of the animals treated intra-articularly with Legend Injectable Solution - 2mL as it did in the saline treated control horses. No gross or histological lesions were observed in the soft tissues or the surface areas of the treated joint.

STORAGE:

Do not store above 40° C (104° F).

HOW SUPPLIED:

Legend Injectable Solution is supplied in a carton of six 2 mL (20 mg) bottles.

Legend Injectable Solution is supplied in a carton of six 4 mL (40 mg) bottles.

Legend Multi Dose Injectable Solution is supplied in 20 mL bottles.

REFERENCE:

¹Swanstrom, O.G. 1978. Hyaluronate (hyaluronic acid) and its use, Proc. American Assoc. Equine Pract., 24th annual convention, pp. 345-348.

For customer service or to obtain product information, including a Material Safety Data Sheet, call 1-800-633-3796.

Legend Multi Dose Injectable Solution	08904327, 08904793, R.1
	September, 2010
Legend Injectable Solution 2 mL -	08710980-017599, 08715516-79001750, R.12
	September, 2010
Legend Injectable Solution 4 mL -	08710999-017699, 08715524-79001760, R.8
	September, 2010

NADA 140-883, Approved by FDA	Made in U.S.A.
15816	© 2010 Bayer HealthCare LLC

Bayer (reg'd), the Bayer Cross (reg'd), Legend® and the horse logo are trademarks of Bayer.

Bayer HealthCare LLC, Animal Health Division, Shawnee Mission, Kansas 66201 U.S.A.

Regulations for product use are established by country. Information contained on this site pertains only to the United States of America, and is not intended to provide adequate information for product use. Before using or dispensing any product, read and carefully observe the label directions.