

INTERVET INC., MERCK ANIMAL HEALTH

556 MORRIS AVE., SUMMIT, NJ, 07901

Telephone: 862-245-4321

Order Desk: 800-648-2118

Fax: 862-245-4935

Customer Service: 800-521-5767

Website: www.intervet.com

PANACUR® HORSE & CATTLE DEWORMER SUSPENSION 10%



Intervet/Merck Animal Health

(fenbendazole)

(100 mg/mL)

RESIDUE WARNINGS:

Do not use in horses intended for food.

Cattle must not be slaughtered for human consumption within 8 days following treatment.

Do not use at 10 mg/kg in dairy cattle. Dose rate of 10 mg/kg is for beef cattle only. Dose rate of 10 mg/kg in dairy cattle could result in violative residues in milk.

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Keep this and all medication out of the reach of children.

DOSAGE:

Horses: 5 mg/kg (2.3 mg/lb.) for the control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*, *Triodontophorus* spp.), small strongyles (*Cyathostomum* spp., *Cylicocyclus* spp., *Cylicostephanus* spp., *Cylicodontophorus* spp.) and pinworms (*Oxyuris equi*). EXAMPLE: 2.3 mL/100 lb.; 23 mL/1000 lb.

10 mg/kg (4.6 mg/lb.) for the control of ascarids (*Parascaris equorum*). Example: (10 mg/kg); 2.3 mL/50 lb.; 23 mL/500 lb.

Beef and Dairy Cattle: 5 mg/kg (2.3 mg/lb.) for the removal and control of:

Lungworm: (*Dictyocaulus viviparus*).

Stomach worm (adults): *Ostertagia ostertagi* (brown stomach worms). **Stomach worm (adults & 4th stage larvae):** *Haemonchus contortus/placei* (barberpole worm), *Trichostrongylus axei* (small stomach worm).

Intestinal worms (adults & 4th stage larvae): *Bunostomum phlebotomum* (hookworm), *Nematodirus helvetianus* (thread-necked intestinal worm), *Cooperia punctata* and *C. oncophora* (small intestinal worm), *Trichostrongylus colubriformis* (bankrupt worm), *Oesophagostomum radiatum* (nodular worm).

Beef Cattle Only - 10 mg/kg (4.6 mg/lb) for the removal and control of:

Stomach worm (4th stage inhibited larvae): *Ostertagia ostertagi* (Type II Ostertagiasis).

Tapeworm: *Moniezia benedeni*.

▶ **Do not use in dairy cattle at 10 mg/kg** ◀

In beef and dairy cattle, the recommended dose of 5 mg/kg is achieved when 2.3 mL of the drug are given for each 100 lb of body weight. In beef cattle only, the recommended dosage of 10 mg/kg for the treatment of Ostertagiasis Type II (inhibited 4th stage larvae) or tapeworm is achieved when 4.6 mL of the drug is given for each 100 lb of body weight.

EXAMPLES: (Horses and Cattle)

Dose (5 mg / kg)	Dose (10 mg / kg)	Animal Weight
2.3 mL	4.6 mL	100 lb
4.6 mL	9.2 mL	200 lb
6.9 mL	13.8 mL	300 lb
9.2 mL	18.4 mL	400 lb
11.5 mL	23.0 mL	500 lb
23.0 mL	46.0 mL	1,000 lb
34.5 mL	69.0 mL	1,500 lb

DIRECTIONS:

Beef and Dairy Cattle and Horses: Determine the proper dose according to estimated body weight. Administer orally by suitable dosing syringe. Insert nozzle of syringe through the interdental space and deposit the drug on the back of the tongue by depressing the plunger. The drug may also be administered by stomach tube. There are no known contraindications to the use of the drug in cattle or horses. For dairy cattle, there is no milk withdrawal period at 5 mg/kg. Panacur® (fenbendazole) Suspension 10% is approved for use concomitantly with an approved form of trichlorfon. Trichlorfon is approved for the treatment of stomach bots

(*Gasterophilus* spp.) in horses. Refer to the manufacturer's label for directions for use and cautions for trichlorfon.

Regular deworming at intervals of six to eight weeks may be required for horses.

Under conditions of continued exposure to parasites, retreatment may be needed after 4-6 weeks.

Manufactured by: DPT Laboratories, San Antonio, TX 78215

Distributed by: **Intervet Inc.**, Millsboro, DE 19966

NADA #s 104-494 and 128-620, Approved by FDA

1,000 mL (33.8 fl.oz.)	697700-B
	697705-B

NAC No.: 10473431