

INTERVET INC., MERCK ANIMAL HEALTH

556 MORRIS AVE., SUMMIT, NJ, 07901

Telephone: 862-245-4321

Order Desk: 800-648-2118

Fax: 862-245-4935

Customer Service: 800-521-5767

Website: www.intervet.com

SAFE-GUARD® CANINE DEWORMER

Intervet/Merck Animal Health

(fenbendazole)

For dogs only, 6 weeks and older.

Drug Facts

Active ingredients (in each dosage unit): Fenbendazole Granules 22.2% (222 mg/g)

Uses: For the treatment and control of:

Tapeworms (*Taenia pisiformis*)

Roundworms (*Toxocara canis*, *Toxascaris leonina*)

Hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*)

Whipworms (*Trichuris vulpis*)

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

HUMAN WARNING: KEEP OUT OF REACH OF CHILDREN.

Directions: The daily dose for Safe-Guard® Canine is 50 mg/kg (22.7 mg/lb) of body weight.

You should weigh your dog to make sure you are using the right dose; i.e. correct gram amount and appropriate number of pouches.

Your dog must be treated with the following dose for 3 days in a row.

Dog Weight (lbs)	Pouch Size (Daily Dose)
10	1 gram
11 to 20	2 gram
21 to 30*	1 gram + 2 gram
31 to 40	4 gram
41 to 50*	1 gram + 4 gram
51 to 60*	2 gram + 4 gram
61 to 80*	Two 4 gram
Over 80*	* Combine pouches to obtain recommended daily dosage.

Mix daily dose with a small amount of the usual food. Make sure your dog eats all the medicated food. You may need to moisten dry dog food to aid mixing.

Repeat the dose for 3 days in a row.

Recommended Deworming Schedule: Schedule may vary depending on the climate where you live and the activity of your dog. The following schedule should be used as a general guidance.

- Weaned pups (6-8 weeks of age) should be dewormed at 6, 8, 10 and 12 weeks of age.
- Treat the dam at the same time as the puppies.
- Dogs over 6 months of age should be dewormed at least twice each year.

Diagnosis of Parasites: Consult your veterinarian for help in the diagnosis, treatment and control of parasitism. Specific diagnoses require laboratory testing. Dogs can become infected with several different kinds of tapeworms. Safe-Guard[®] Canine only kills *Taenia* species of tapeworms. If you continue to see tapeworm segments in your dog's stool after treatment with Safe-Guard[®] Canine, consult your veterinarian.

Roundworms and hookworms may pose serious human health risks; regular deworming may help reduce those risks.

Product Use: Safe-Guard[®] Canine is safe for use in puppies 6 weeks or older and adult dogs, including pregnant bitches. Do not deworm a dog or puppy that is sick. Consult your veterinarian for diagnosis of the illness.

Adverse Reactions: In US clinical studies 3 of 240 dogs (about 1% of the treated dogs) had vomiting associated with use of the product.

Storage: Store at controlled room temperature (59-86°F).

Questions? Comments?

To obtain product information including material safety data sheets (MSDS) or to report a suspected adverse reaction, call **1-800-441-8272**.

Distributed by: INTERVET INC., Millsboro, DE

Made in Canada

www.Safe-Guard-For-Dogs.com

NADA # 121-473, Approved by FDA

Safe-Guard is a registered trademark of Intervet Inc.

Net Weight:		
3 grams	Three 1 gram pouches for a 3-day treatment. Each pouch treats 10 lbs.	446100-C IC-30205/G 749100-C PF-30207/G
6 grams	Three 2 gram pouches for a 3-day treatment. Each pouch treats 20 lbs.	446200-C IC-30213/G 749200-C PF-30216/G
12 grams	Three 4 gram pouches for a 3-day treatment. Each pouch treats 40 lbs.	446400-C IC-30218/G 749400-C PF-30219/G

NAC No.: 10474291